

INDETERMINACY OF THE MOMENT PROBLEM FOR SYMMETRIC PROBABILITY MEASURES

HAYATO SAIGO AND HIROKI SAKO

ABSTRACT. In this paper, the moment problem for symmetric probability measures is characterized in terms of associated sequences called Jacobi sequences $\{\omega_n\}$. A notion named property (SC), which is proved to be a necessary and sufficient condition for the indeterminacy of the moment problem, naturally arises from the viewpoint of finite dimensional approximation for infinite matrices. We prove that the moment problem for q -Gaussian not only for $q > 1$ but also for $q < -1$ is indeterminate. We also prove that hyperbolic secant distribution is “the last probability measure” which is uniquely determined by the moment sequence of power type, just by checking property (SC) with quite easy calculation.

1. INTRODUCTION

The moment sequence of a probability measure is, if it exists, one of the most important information of the measure. Then, does the moment sequence determine the measure uniquely or not?—This fundamental question called “moment problem” (or more precisely “determinate moment problem”) has been attracted many mathematicians since the work by Stieltjes [Sti94], who proposed and solved the moment problem for $\mathbb{R}_{\geq 0}$. It was Hamburger ([Ham20a], [Ham20b], and [Ham21]) who extended the problem for $\mathbb{R}$, on which we study here. (For the subject and its history, see Akhiezer [Akh65] and Shohat–Tamarkin [ST43]). If the measure is unique, the problem is said to be determinate, and otherwise, indeterminate. In the present paper, we discuss the moment problem in terms of the associated sequence which we call “Jacobi sequence”.

The sequence of orthogonal polynomials also provides useful data for the probability measure μ whose moments $\{M_n = \int_{\mathbb{R}} x^n d\mu\}$ are finite. Then the space of polynomial functions is contained in the Hilbert space $L^2(\mathbb{R}, \mu)$. A Gram-Schmidt procedure provides orthogonal polynomials which only depend on the moment sequence. Let $\{P_n(x)\}_{n=0,1,\dots}$ be the monic orthogonal polynomials of μ . Then there exist sequences $\{\alpha_n\}_{n=0,1,\dots}$ and $\{\omega_n\}_{n=1,2,\dots}$ such that

$$(1) \quad xP_n(x) = P_{n+1}(x) + \alpha_{n+1}P_n(x) + \omega_n P_{n-1}(x), \quad P_{-1}(x) = 0.$$

These sequences $\{\omega_n\}$ and $\{\alpha_n\}$ are called “Jacobi sequences” associated to μ . The Jacobi sequences have the same amount of data as the moment sequence.

In this paper, we will concentrate on the case that μ is symmetric, i.e., $\mu(-dx) = \mu(dx)$. Then α_n are all zero. We use the term “Jacobi sequence” to indicate the

2000 *Mathematics Subject Classification.* 47A57, 44A60.

Key words and phrases. Probability Measure, Moment Problem, Jacobi Sequence.

sequence $\{\omega_n\}$. It is also known that the real numbers ω_n are all greater than 0 if and only if μ has infinite support. Since the moment problem is trivial for measures with finite support, we always assume that $\{\omega_n\}$ are all positive.

In the present paper, a new notion “property (SC)”¹ is defined for infinite Jacobi sequences and is proved to be equivalent to the indeterminacy of the moment problem. Section 2 is devoted to formulate the notion. In Section 3, it is shown that property (SC) leads to a construction of two distinct self-adjoint operators and proved to imply the indeterminacy of the moment problem. We will investigate basic examples of Jacobi sequences with property (SC), q -deformed integers $[n]_q = 1 + q + q^2 + \cdots + q^{n-1}$ ($q > -1$), $(-1)^{n-1} \frac{q^n - 1}{q - 1}$ ($q < -1$) and powers of integers n^p ($p > 2$) in Section 4. The following theorems are consequences:

Theorem 1.1. *The moment problem for q -Gaussians ($q \in \mathbb{R}$) are indeterminate if and only if $|q| > 1$.*

In the case of $q > 1$, Ismail and Masson proved that the moment problem is indeterminate, identifying extremal measures [IM94].

Theorem 1.2. *Let p be a positive number. The probability measure corresponding to the Jacobi sequence $\{n^p\}$ is unique if and only if $p \leq 2$.*

The Jacobi sequence of the the hyperbolic secant distribution $\frac{dx}{e^{\pi x/2} + e^{-\pi x/2}}$ on $\mathbb{R}$ is $\{n^2\}_{n=1}^\infty$. The theorem above means that the hyperbolic secant distribution is “the last probability measure” which is uniquely determined by the moment sequence of power type. In Section 5, we prove that property (SC) holds if and only if the corresponding probability measure is not unique.

Notations and Conventions.

- $\mathbb{N}$ is the set of non-negative integers $\{0, 1, 2, \dots\}$.
- $\ell_2(\mathbb{N})$ stands for the set of all the square summable sequences.
- Identify the finite dimensional Hilbert space $\ell_2(\{0, 1, \dots, m\})$ with the closed subspace of $\ell_2(\mathbb{N})$.
- We regard a vector η in $\ell_2(\{0, 1, \dots, m\})$ or in $\ell_2(\mathbb{N})$ as a column vector. For an natural number l , ${}^l\eta$ stands for the l -th entry of the vector.
- For $k \in \mathbb{N}$, δ_k stands for a unit vector given by ${}^l\delta_k = \begin{cases} 0, & l \neq k, \\ 1, & l = k. \end{cases}$
- Fixing the orthonormal basis $\{\delta_k\}$, we identify the set of linear operators $\mathbb{B}(\ell_2(\{0, 1, \dots, m\}))$ with the set of complex matrices $\mathbb{M}(m+1)$.

¹ Since the announcement of this result, many experts have expressed their impression that property (SC) has been already discovered. Of course, in a logical sense, there exist many conditions which are equivalent to property (SC), because the determinate moment problem is classical. The problem has been rephrased in terms of (absolute) convergence or divergence of several kinds of sequences. However, the authors did not find similarity between known conditions and our property (SC). Our motivation for the definition of property (SC) is to give a concise and self-contained proof for the indeterminacy for interesting measures in quantum probability theory. As an example, our characterization implies the new theorems (Theorem 1.1 and Theorem 1.2).

- For an operator $Y \in \mathbb{B}(\ell_2(\{0, 1, \dots, m\}))$, ${}^l Y_k$ is the coefficient $\langle Y \delta_k, \delta_l \rangle$.
- We write Y_k for the k -th column vector $({}^l Y_k)_{l=0}^m$ of Y .
- We use the same notations for an operator $Y \in \mathbb{B}(\ell_2(\mathbb{N}))$. That is, ${}^l Y_k$ stands for $\langle Y \delta_k, \delta_l \rangle$ and Y_k stands for $({}^l Y_k)_{l=0}^\infty \in \ell_2(\mathbb{N})$.
- $\frac{a_1}{b_1} + \frac{a_2}{b_2} + \dots + \frac{a_m}{b_m}$ is the continued fraction
$$\frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \frac{a_4}{b_4 + \frac{a_5}{b_5 + \frac{a_6}{b_6 + \frac{a_7}{b_7 + \frac{a_8}{b_8 + \frac{a_9}{b_9 + \frac{a_{10}}{b_{10}}}}}}}}}}}}$$

2. FLUCTUATION PROPERTY OF THE STIELTJES TRANSFORM AT i

The three-term recurrence relation (equation (1)) give a bijective correspondence between the moment sequences of symmetric probability measures

$$\{\{M_n\}_{n=0}^\infty; \exists \text{ prob. } \mu, d\mu(x) = d\mu(-x), M_n = \int_{\mathbb{R}} x^n d\mu, \#(\text{supp } \mu) = \infty\},$$

and the infinite Jacobi sequences $\{\{\omega_n\}_{n=1}^\infty; \omega_n > 0\}$. For an infinite Jacobi sequence $\{\omega_n\}_{n=1}^\infty$, denote by X the infinite matrix corresponding to the sequence:

$$X = \begin{pmatrix} 0 & \sqrt{\omega_1} & 0 & \cdots \\ \sqrt{\omega_1} & 0 & \sqrt{\omega_2} & \\ 0 & \sqrt{\omega_2} & 0 & \ddots \\ \vdots & & \ddots & \ddots \end{pmatrix}.$$

We regard X as an operator defined on $\oplus_{\mathbb{N}} \mathbb{C} \subset \ell_2(\mathbb{N})$. The operator X is a densely defined symmetric operator on the Hilbert space $\ell_2(\mathbb{N})$. We note that X is not essentially self-adjoint in general and that X^* is not necessarily self-adjoint. In the case that the growth rate of the Jacobi sequence $\{\omega_n\}$ is large, there exist more than one self-adjoint operators between X and X^* . We show the existence of these operators, by exploiting finite dimensional approximations of the operator X .

Let X_m denote the $(m+1) \times (m+1)$ matrix

$$X_m = \begin{pmatrix} 0 & \sqrt{\omega_1} & 0 & \cdots & 0 \\ \sqrt{\omega_1} & 0 & \sqrt{\omega_2} & \ddots & \vdots \\ 0 & \sqrt{\omega_2} & 0 & \ddots & 0 \\ \vdots & \ddots & \ddots & \ddots & \sqrt{\omega_m} \\ 0 & \cdots & 0 & \sqrt{\omega_m} & 0 \end{pmatrix}$$

corresponding to the finite Jacobi sequence $\{\omega_1, \omega_2, \dots, \omega_m\}$. We regard X_m as an operator acting on the Hilbert space $\ell_2(\{0, 1, \dots, m\})$.

We begin with investigating the Stieltjes transform $G_m(i)$ of X_m at i with respect to the state $\langle \cdot, \delta_0 \rangle$. The complex number $G_m(i)$ is equal to the upper-left matrix

coefficient of $(i - X_m)^{-1}$. This quantity is expressed by the continued fraction ²

$$\frac{1}{i} - \frac{\omega_1}{i} - \frac{\omega_2}{i} - \cdots - \frac{\omega_m}{i}.$$

The quantity $iG_m(i)$ is simply expressed as $\frac{1}{1} + \frac{\omega_1}{1} + \frac{\omega_2}{1} + \cdots + \frac{\omega_m}{1}$. To study $\{iG_m(i)\}_{m=1}^\infty$, we define sequences $\{A_m\}_{m=-1}^\infty$ and $\{B_m\}_{m=-1}^\infty$ by

$$\begin{aligned} A_{-1} &= 0, & A_0 &= 1, & A_m &= A_{m-1} + \omega_m A_{m-2}, \\ B_{-1} &= 1, & B_0 &= 1, & B_m &= B_{m-1} + \omega_m B_{m-2}. \end{aligned}$$

Lemma 2.1. *The Stieltjes transform $G_m(i)$ of X_m is expressed by*

$$iG_m(i) = \frac{A_m}{B_m}.$$

Proof. Define A_{-2} and B_{-2} by $A_{-2} = 1$, $B_{-2} = 0$. Define ω_0 by 1. Applying the equation (1.71) of Hora–Obata [HO07], we obtain the proposition. $\square$

Lemma 2.2. *For $m \geq 1$, $A_m B_{m-1} - B_m A_{m-1} = (-1)^m \omega_1 \omega_2 \cdots \omega_m$.*

Proof. We prove by induction. For the case of $m = 1$, we have

$$A_1 B_0 - B_1 A_0 = 1 \cdot 1 - (1 + \omega_1) \cdot 1 = -\omega_1.$$

Assume that the lemma holds. Then we have

$$\begin{aligned} A_{m+1} B_m - A_m B_{m+1} &= (A_m + \omega_{m+1} A_{m-1}) B_m - A_m (B_m + \omega_{m+1} B_{m-1}) \\ &= -\omega_{m+1} (A_m B_{m-1} - B_m A_{m-1}) \\ &= (-1)^{m+1} \omega_1 \cdots \omega_m \omega_{m+1}. \end{aligned}$$

$\square$

Lemma 2.3. *For $m \geq 2$, $iG_m(i) - iG_{m-1}(i) = \frac{(-1)^m \omega_1 \omega_2 \cdots \omega_m}{B_{m-1} B_m}$*

Proof. By Lemma 2.1, we have

$$iG_m(i) - iG_{m-1}(i) = \frac{A_m}{B_m} - \frac{A_{m-1}}{B_{m-1}} = \frac{A_m B_{m-1} - A_{m-1} B_m}{B_m B_{m-1}}.$$

Lemma 2.2 yields the desired equation. $\square$

It follows that the sequence $\{iG_m(i)\}_{m=1}^\infty$ alternately increases and decreases. More precisely,

- If m is odd, then $iG_{m-1}(i) > iG_m(i)$;
- If m is even, then $iG_{m-1}(i) < iG_m(i)$.

We next investigate the difference $|iG_m(i) - iG_{m-1}(i)|$.

Proposition 2.4. *For $m \geq 3$, $\frac{|iG_m(i) - iG_{m-1}(i)|}{|iG_{m-1}(i) - iG_{m-2}(i)|} = 1 - \frac{B_{m-1}}{B_m} < 1$.*

²This equality has been shown in many references. See, e.g., Hora–Obata [HO07, Lemma 1.87]. This is also shown in the first equation of Lemma 3.3.

Proof. By Lemma 2.3, we have

$$\frac{|iG_m(i) - iG_{m-1}(i)|}{|iG_{m-1}(i) - iG_{m-2}(i)|} = \frac{\omega_1 \cdots \omega_m \frac{B_{m-2}B_{m-1}}{\omega_1 \cdots \omega_{m-1}}}{\frac{B_{m-1}B_m}{\omega_1 \cdots \omega_{m-1}}} = \left| \frac{\omega_m B_{m-2}}{B_m} \right|$$

By the definition of B_m , the above quantity is equal to $(B_m - B_{m-1})/B_m$. $\square$

The above proposition implies the following properties of $\{iG_m(i)\}_{m=1}^\infty$.

Theorem 2.5.

- The subsequence $\{iG_m(i); m \text{ even}\}$ strictly decreases.
- The subsequence $\{iG_m(i); m \text{ odd}\}$ strictly increases.
- If n is an even natural number, $iG_n(i)$ is an upper bound of $\{iG_m(i); m \text{ odd}\}$.
- If n is an odd natural number, $iG_n(i)$ is a lower bound of $\{iG_m(i); m \text{ even}\}$.

Define two complex numbers G_{even} and G_{odd} by

$$iG_{\text{even}} = \lim_{m \rightarrow \infty} iG_{2m}(i), \quad iG_{\text{odd}} = \lim_{m \rightarrow \infty} iG_{2m+1}(i).$$

Proposition 2.6. *The following conditions are equivalent:*

- (1) $G_{\text{even}} \neq G_{\text{odd}}$;
- (2) $\lim_{m \rightarrow \infty} \prod_{n=1}^m \left(1 - \frac{B_{n-1}}{B_n}\right) > 0$;
- (3) $\lim_{m \rightarrow \infty} \sum_{n=1}^m \frac{B_{n-1}}{B_n} < \infty$.

Proof. Since $B_0 = 1$, $B_1 = 1 + \omega_1$ and $B_2 = 1 + \omega_1 + \omega_2$, we have

$$\begin{aligned} iG_2(i) - iG_1(i) &= \frac{1}{1 + \omega_1/(1 + \omega_2)} - \frac{1}{1 + \omega_1} \\ &= \frac{\omega_1}{1 + \omega_1} \frac{\omega_2}{1 + \omega_1 + \omega_2} \\ &= \left(1 - \frac{B_0}{B_1}\right) \left(1 - \frac{B_1}{B_2}\right). \end{aligned}$$

Equivalence between (1) and (2) follows from the equality

$$|iG_m(i) - iG_{m-1}(i)| = (iG_2(i) - iG_1(i)) \prod_{n=3}^m \left(1 - \frac{B_{n-1}}{B_n}\right) = \prod_{n=1}^m \left(1 - \frac{B_{n-1}}{B_n}\right).$$

Equivalence between (2) and (3) is well-known. ³ $\square$

Definition 2.7. *If one of the equivalent conditions in Proposition 2.6 holds, then the infinite Jacobi sequence $\{\omega_m\}_{m=1}^\infty$ is said to have property (SC). ⁴*

³Use the inequality $-2x < \log(1 - x) < -x$, $0 < x \leq 1/2$.

⁴Property (SC) means that the subsequences of the Stieltjes transform ‘Separately Converge.’

$iG_m(i)$ is in the closed interval

$$[iG_1(i), iG_2(i)] = \left[\frac{1}{1} + \frac{\omega_1}{1}, \frac{1}{1} + \frac{\omega_1}{1} + \frac{\omega_2}{1} \right].$$

It follows that $D_0^{(m)}$ is in the closed interval $\left[\frac{\omega_1}{1 + \omega_2}, \omega_1 \right]$. By Theorem 2.5, the sequences $\{D_0^{(m)}; m \text{ is even}\}$ and $\{D_0^{(m)}; m \text{ is odd}\}$ converge to real numbers.

Suppose that the lemma holds for a natural number n . The sequence $\{D_{n+1}^{(m)}\}_{m=n+2}^\infty$ is described as $D_{n+1}^{(m)} = \omega_{n+1}/D_n^{(m)} - 1$. Since

$$D_n^{(m)} = \frac{\omega_{n+1}}{1} + \frac{\omega_{n+2}}{1} + \cdots + \frac{\omega_m}{1}$$

is in the closed interval $\left[\frac{\omega_{n+1}}{1} + \frac{\omega_{n+2}}{1}, \frac{\omega_{n+1}}{1} + \frac{\omega_{n+2}}{1} + \frac{\omega_{n+3}}{1} \right]$, we have

$$D_{n+1}^{(m)} = \frac{\omega_{n+1}}{D_n^{(m)}} - 1 \in \left[\frac{\omega_{n+2}}{1 + \omega_{n+3}}, \omega_{n+2} \right].$$

By the hypothesis of induction, the above lemma for $n + 1$ follows. $\square$

We denote by $Y^{(m)}$ the matrix $(i - X_m)^{-1} \in \mathbb{B}(\ell_2(\{0, 1, \dots, m\}))$. We decompose $Y^{(m)}$ into column vectors as $Y^{(m)} = [Y_0^{(m)} Y_1^{(m)} \cdots Y_m^{(m)}]$. We describe the vector $Y_0^{(m)}$

as $Y_0^{(m)} = \begin{pmatrix} {}^0Y_0^{(m)} \\ {}^1Y_0^{(m)} \\ \vdots \\ {}^mY_0^{(m)} \end{pmatrix}$. The vector $Y_0^{(m)}$ is described by C_n and $D_n^{(m)}$ as follows.

Lemma 3.3.

$$\begin{aligned} {}^0Y_0^{(m)} &= \frac{1}{i} \frac{1}{1 + D_0^{(m)}}, \\ {}^nY_0^{(m)} &= \frac{\sqrt{\omega_1 \cdots \omega_n} C_1 \cdots C_n}{i^{n+1}} \frac{1}{1 + C_n D_n^{(m)}}, \quad 1 \leq n \leq m-1, \\ {}^mY_0^{(m)} &= \frac{\sqrt{\omega_1 \cdots \omega_m} C_1 \cdots C_m}{i^{m+1}}. \end{aligned}$$

Proof. We define a vector $\tilde{Y} = \begin{pmatrix} {}^0\tilde{Y} \\ \vdots \\ {}^m\tilde{Y} \end{pmatrix}$ by the right hand side of the equations which we are going to prove. By the injectivity of $(i - X_m)^{-1}$, it suffices to show

that $\begin{pmatrix} {}^0Z \\ \vdots \\ {}^mZ \end{pmatrix} = (i - X_m)\tilde{Y}$ is equal to $\delta_0 = \begin{pmatrix} 1 \\ 0 \\ \vdots \end{pmatrix}$. Equation (2) and (3) yield

$$\begin{aligned} {}^0Z &= i \left({}^0\tilde{Y} \right) - \sqrt{\omega_1} \left({}^1\tilde{Y} \right) \\ &= \frac{1}{1 + D_0^{(m)}} + \frac{\omega_1 C_1}{1 + C_1 D_1^{(m)}} \\ &= \frac{1 + D_1^{(m)}}{1 + D_1^{(m)} + \omega_1} + \frac{\omega_1}{C_1^{-1} + D_1^{(m)}} \\ &= 1. \end{aligned}$$

In order to prove ${}^nZ = 0, 1 \leq n \leq m-1$, we compute

$$\begin{aligned} & -\sqrt{\omega_n}({}^{n-1}\tilde{Y}) + i({}^n\tilde{Y}) \\ &= \frac{\sqrt{\omega_1 \cdots \omega_n} C_1 \cdots C_{n-1}}{i^{n+2}} \left(\frac{1}{1 + C_{n-1} D_{n-1}^{(m)}} - \frac{C_n}{1 + C_n D_n^{(m)}} \right). \end{aligned}$$

By $D_{n-1}^{(m)} = \omega_n / (1 + D_n^{(m)})$ (equation (3)) and $1 + \omega_n C_{n-1} = C_n^{-1}$ (equation (2)), we have

$$\begin{aligned} \frac{1}{1 + C_{n-1} D_{n-1}^{(m)}} - \frac{C_n}{1 + C_n D_n^{(m)}} &= \frac{1 + D_n^{(m)}}{1 + \omega_n C_{n-1} + D_n^{(m)}} - \frac{1}{C_n^{-1} + D_n^{(m)}} \\ &= \frac{D_n^{(m)}}{C_n^{-1} + D_n^{(m)}} \\ &= C_n \frac{1}{1/D_n^{(m)} + C_n} \\ &= \omega_{n+1} C_n \frac{1}{1 + \omega_{n+1} C_n + D_{n+1}^{(m)}}. \end{aligned}$$

Equation (2) for $n+1$ yields

$$\frac{1}{1 + \omega_{n+1} C_n + D_{n+1}^{(m)}} = \frac{1}{1/C_{n+1} + D_{n+1}^{(m)}} = \frac{C_{n+1}}{1 + C_{n+1} D_{n+1}^{(m)}}.$$

It follows that

$$\begin{aligned} {}^nZ &= -\sqrt{\omega_n} \left({}^{n-1}\tilde{Y} \right) + i \left({}^n\tilde{Y} \right) - \sqrt{\omega_{n+1}} \left({}^{n+1}\tilde{Y} \right) \\ &= \frac{\sqrt{\omega_1 \cdots \omega_n} C_1 \cdots C_{n-1}}{i^{n+2}} \omega_{n+1} C_n \frac{C_{n+1}}{1 + C_{n+1} D_{n+1}^{(m)}} - \sqrt{\omega_{n+1}} \left({}^{n+1}\tilde{Y} \right) \\ &= 0. \end{aligned}$$

Since $D_{m-1}^{(m)}$ is equal to ω_m , we compute ${}^mZ$ as follows:

$$\begin{aligned} {}^mZ &= -\sqrt{\omega_m} \left({}^{m-1}\tilde{Y} \right) + i \left({}^m\tilde{Y} \right) \\ &= \frac{\sqrt{\omega_1 \cdots \omega_m} C_1 \cdots C_{m-1}}{i^m} \left(\frac{-1}{1 + C_{m-1} D_{m-1}^{(m)}} + C_m \right) \\ &= 0 \end{aligned}$$

It follows that $(i - X_m)\tilde{Y}$ is equal to δ_0 . $\square$

We regard $Y^{(m)} = (i - X_m)^{-1} \in \mathbb{B}(\ell_2(\{0, 1, \dots, m\}))$ as an operator in $\mathbb{B}(\ell_2(\mathbb{N}))$.

Lemma 3.4. *The sequences $\{Y^{(2m)}\}_{m=0}^\infty$ and $\{Y^{(2m+1)}\}_{m=0}^\infty$ converge in the weak operator topology.*

Recall that ${}^lY_k^{(m)}$ stands for the (l, k) -entry of the matrix $Y^{(m)}$.

Proof. Note that the operator norm of $Y^{(m)}$ is at most 1, because X_m is self-adjoint and $Y^{(m)}$ is the resolvent at i . It suffices to show that for arbitrary natural numbers $k, l \in \mathbb{N}$, the sequences $\{{}^lY_k^{(2m)}\}_{m=0}^\infty$ and $\{{}^lY_k^{(2m+1)}\}_{m=0}^\infty$ converge. We prove this claim by induction on k .

By the concrete expressions of ${}^lY_0^{(2m)}$ and ${}^lY_0^{(2m+1)}$ in Lemma 3.3 and by Lemma 3.2, $\{{}^lY_0^{(2m)}\}_m$ and $\{{}^lY_0^{(2m+1)}\}_m$ respectively converge. By the equation $Y^{(m)}(i - X_m) = \text{id}_{\ell_2(\{0, 1, \dots, m\})}$, we have $iY_0^{(m)} - \sqrt{\omega_1}Y_1^{(m)} = \delta_0$. Therefore the column vector $Y_1^{(m)}$ is equal to $(iY_0^{(m)} - \delta_0)/\sqrt{\omega_1}$ and their entries converge.

Assume that the above claim holds for the column vectors $Y_{k-2}^{(m)}$ and $Y_{k-1}^{(m)}$. For the case of $2 \leq k < m$, observing the $(k-1)$ -st column of the equation $Y^{(m)}(i - X_m) = \text{id}_{\ell_2(\{0, 1, \dots, m\})}$, we have

$$\begin{aligned} -\sqrt{\omega_{k-1}}Y_{k-2}^{(m)} + iY_{k-1}^{(m)} - \sqrt{\omega_k}Y_k^{(m)} &= \delta_{k-1}, \\ Y_k^{(m)} &= \frac{-\sqrt{\omega_{k-1}}Y_{k-2}^{(m)} + iY_{k-1}^{(m)} - \delta_{k-1}}{\sqrt{\omega_k}}. \end{aligned}$$

By the assumption, $\{{}^kY_l^{(2m)}\}_m$ and $\{{}^kY_l^{(2m+1)}\}_m$ respectively converge. $\square$

In the case that property (SC) holds, we obtain a stronger conclusion in Proposition 3.7. Define an infinite sequence $\hat{Y} = \begin{pmatrix} {}^0\hat{Y} \\ {}^1\hat{Y} \\ \vdots \end{pmatrix}$ by

$${}^0\hat{Y} = 1, {}^1\hat{Y} = \sqrt{\omega_1}, {}^m\hat{Y} = \sqrt{\omega_1 \cdots \omega_m} C_1 \cdots C_m.$$

Lemma 3.5. *The sequence $\hat{Y}$ is an element of $\ell_2(\mathbb{N})$, if property (SC) holds.*

Proof. By the inequality $\omega_{n+1}C_nC_{n+1} = \omega_{n+1}C_n \frac{1}{1 + \omega_{n+1}C_n} \leq 1$, we estimate $({}^m\widehat{Y})^2$ by

$$({}^m\widehat{Y})^2 = \omega_1 C_1 \left(\prod_{n=1}^{m-1} \omega_{n+1} C_n C_{n+1} \right) C_m \leq \omega_1 C_1 C_m.$$

If $\{\omega_m\}_{m=1}^\infty$ has property (SC), then $\|\widehat{Y}\|_2^2 \leq 1 + \omega_1 C_1 \sum_{m=1}^\infty C_m < \infty$. $\square$

Lemma 3.6. *If property (SC) holds, then there exist positive square summable sequences $\{\widehat{Y}_k = ({}^l\widehat{Y}_k)_{l=0}^\infty; k \in \mathbb{N}\} \subset \ell_2(\mathbb{N})$ such that for every $k, l \in \mathbb{N}$ and $m \geq k$, $|{}^l Y_k^{(m)}| \leq {}^l\widehat{Y}_k$.*

Proof. We prove by induction on k . Define $\widehat{Y}_0$ by $\widehat{Y}$. By Lemma 3.3 and the definition of $\widehat{Y}$, we have $|{}^l Y_0^{(m)}| \leq {}^l\widehat{Y}_0$. For $m \geq 1$, the vector $Y_1^{(m)}$ is expressed by $(iY_0^{(m)} - \delta_0)/\sqrt{\omega_1}$. Absolute values of entries are dominated by those of $(\widehat{Y}_0 + \delta_0)/\sqrt{\omega_1}$. Define $\widehat{Y}_1$ by $(\widehat{Y}_0 + \delta_0)/\sqrt{\omega_1}$.

For $m \geq k \geq 1$, since the $(k-1)$ -st column of $Y^{(m)}(i - X_m)$ is

$$\delta_{k-1} = -\sqrt{\omega_{k-1}}Y_{k-2}^{(m)} + iY_{k-1}^{(m)} - \sqrt{\omega_k}Y_k^{(m)},$$

the column vector $Y_k^{(m)}$ is expressed by $(iY_{k-1}^{(m)} - \sqrt{\omega_{k-1}}Y_{k-2}^{(m)} - \delta_{k-1})/\sqrt{\omega_k}$. Absolute values of entries are dominated by those of

$$\widehat{Y}_k = \frac{\widehat{Y}_{k-1} + \sqrt{\omega_{k-1}}\widehat{Y}_{k-2} + \delta_{k-1}}{\sqrt{\omega_k}}.$$

Repeating this procedure, we obtain vectors $\{\widehat{Y}_k\}$ satisfying the above lemma. $\square$

Proposition 3.7. *If property (SC) holds, then the sequences $\{Y^{(2m)}\}_m \subset \mathbb{B}(\ell_2(\mathbb{N}))$ and $\{Y^{(2m+1)}\}_m \subset \mathbb{B}(\ell_2(\mathbb{N}))$ respectively converge in the strong operator topology.*⁵

Proof. Suppose that property (SC) holds. For every $k \in \mathbb{N}$, the sequence of the k -th column vectors $\{Y_k^{(2m)}\}_m$ strongly converges. Indeed, by Lemma 3.4, for every $l \in \mathbb{N}$, $\{{}^l Y_k^{(2m)}\}_m$ converges. For $m \geq k/2$, the absolute value of $Y_k^{(2m)}$ dominated by a square summable positive sequence $\widehat{Y}_k$. By Lebesgue's dominated convergence theorem, $\{Y_k^{(2m)}\}_m$ strongly converges. By the same proof, for every k , the sequence of vectors $\{Y_k^{(2m+1)}\}_m$ also strongly converges.

Since the operator norms of $Y^{(m)} = (i - X_m)^{-1}$ are at most 1, the sequences $\{Y^{(2m)}\}_{m=0}^\infty$ and $\{Y^{(2m+1)}\}_{m=0}^\infty$ respectively converge in the strong operator topology. $\square$

Now we are ready to construct two self-adjoint operators X_{even} and X_{odd} , in the case that $\{\omega_n\}$ holds property (SC). Let Y^{even} denote the strong limit of $\{Y^{(2m)}\}_{m=0}^\infty$ and let Y^{odd} denote the strong limit of $\{Y^{(2m+1)}\}_{m=0}^\infty$. Recall that X^* is the adjoint of the Jacobi matrix $X : \oplus_{\mathbb{N}} \mathbb{C} \rightarrow \oplus_{\mathbb{N}} \mathbb{C} \subset \ell_2(\mathbb{N})$.

⁵Property (SC) is a 'Sufficient Condition' for the 'Strong Convergence.'

Lemma 3.8. *If property (SC) holds, then*

$$\begin{aligned} \text{Image}(Y^{\text{even}}) &\subset \text{Dom}(X^*), & (i - X^*)Y^{\text{even}} &= \text{id}_{\ell_2(\mathbb{N})}, \\ \text{Image}(Y^{\text{odd}}) &\subset \text{Dom}(X^*), & (i - X^*)Y^{\text{odd}} &= \text{id}_{\ell_2(\mathbb{N})}. \end{aligned}$$

Proof. We prove the first equality. It suffices to show

$$\langle Y^{\text{even}}\xi, (-i - X)\eta \rangle = \langle \xi, \eta \rangle,$$

for $\xi \in \text{Image}(Y^{\text{even}})$, and every finitely supported vector $\eta \in \oplus_{\mathbb{N}} \mathbb{C}$. If m is so large that $\{0, 1, \dots, 2m-1\}$ includes $\text{supp}(\eta)$, then

$$\langle Y^{\text{even}}\xi, (-i - X)\eta \rangle = \langle Y^{\text{even}}\xi, (-i - X_{2m})\eta \rangle.$$

Therefore we have

$$\begin{aligned} \langle Y^{\text{even}}\xi, (-i - X)\eta \rangle &= \lim_m \langle Y^{(2m)}\xi, (-i - X_{2m})\eta \rangle \\ &= \lim_m \langle (i - X_{2m})Y^{(2m)}\xi, \eta \rangle \\ &= \lim_m \langle \xi, P_{2m}\eta \rangle, \end{aligned}$$

where P_{2m} stands for the orthogonal projection onto $\ell_2(\{0, 1, \dots, 2m\})$. It follows that $\langle Y^{\text{even}}\xi, (-i - X)\eta \rangle = \langle \xi, \eta \rangle$. $\square$

Lemma 3.9. *If property (SC) holds, then*

$$\text{Image}(Y^{\text{even}})^* = \text{Image}(Y^{\text{even}}), \quad \text{Image}(Y^{\text{odd}})^* = \text{Image}(Y^{\text{odd}}).$$

Proof. Since the matrix coefficients of $i - X_m$ are symmetric, those of $Y^{(m)}$ are also symmetric. More precisely, ${}_k Y_l^{(m)}$ is equal to ${}_l Y_k^{(m)}$. It follows that the adjoint $(Y^{(m)})^*$ is equal to $\overline{Y^{(m)}} = \left(\overline{{}_l Y_k^{(m)}} \right)_{l,k}$.

Notice that all the arguments in this paper is also valid, even if we replace ‘ i ’ with ‘ $-i$.’ It follows that the sequences

$$\{(Y^{(2m)})^* = \overline{Y^{(2m)}}\}_{m=0}^\infty, \quad \{(Y^{(2m+1)})^* = \overline{Y^{(2m+1)}}\}_{m=0}^\infty$$

strongly converge. Observing their matrix coefficients, we have

$$\lim_m (Y^{(2m)})^* = \overline{Y^{\text{even}}} = (Y^{\text{even}})^*, \quad \lim_m (Y^{(2m+1)})^* = \overline{Y^{\text{odd}}} = (Y^{\text{odd}})^*.$$

By the resolvent identity, we have

$$\begin{aligned} (-i - X_{2m})^{-1} &= (i - X_{2m})^{-1} + 2i(i - X_{2m})^{-1}(-i - X_{2m})^{-1}, \\ (Y^{(2m)})^* &= Y^{(2m)} \left(1 + 2i\overline{Y^{(2m)}} \right) = Y^{(2m)}(1 + 2i(Y^{(2m)})^*). \end{aligned}$$

Taking limits with respect to the strong operator topology, we have

$$(Y^{\text{even}})^* = Y^{\text{even}}(1 + 2i(Y^{\text{even}})^*).$$

It follows that the image of $(Y^{\text{even}})^*$ is included in that of Y^{even} . Replacing ‘ i ’ with ‘ $-i$,’ we have

$$\text{Image}(Y^{\text{even}})^* = \text{Image}Y^{\text{even}}.$$

By the same proof, we also obtain the equality $\text{Image}(Y^{\text{odd}})^* = \text{Image}Y^{\text{odd}}$. $\square$

Lemma 3.10. *Let Y be a bounded operator on a Hilbert space $\mathcal{H}$. Suppose that Y and its adjoint Y^* are injective. Then the images of Y , Y^* are dense in $\mathcal{H}$. The adjoint of $Y^{-1}: \text{Image}(Y) \rightarrow \mathcal{H}$ is equal to the inverse of Y^* .*

Proof. By the equalities $\overline{\text{Image}(Y)} = \text{Ker}(Y^*)^\perp$ and $\overline{\text{Image}(Y^*)} = \text{Ker}(Y)^\perp$, we obtain the first assertion.

For a closed operator Z on $\mathcal{H}$, let $\mathcal{G}(Z)$ denote the graph

$$\{\xi \oplus Z\xi; \xi \in \text{Dom}(Z)\} \subset \mathcal{H} \oplus \mathcal{H}.$$

We define two unitary operators U, V on $\mathcal{H} \oplus \mathcal{H}$ by

$$V(\xi \oplus \eta) = (-\eta) \oplus \xi, \quad U(\xi \oplus \eta) = \eta \oplus \xi.$$

Then we have $\mathcal{G}(Z^*) = (V\mathcal{G}(Z))^\perp$. When Z is injective, the graph of

$$Z^{-1}: \text{Image}(Z) \rightarrow \text{Dom}(Z)$$

is given by $\mathcal{G}(Z^{-1}) = U\mathcal{G}(Z)$.

The graph of $(Y^*)^{-1}$ is given by $U((V\mathcal{G}(Y))^\perp)$. The graph of $(Y^{-1})^*$ is given by $(VU\mathcal{G}(Y))^\perp$. Since $-UV = VU$, we obtain $\mathcal{G}((Y^*)^{-1}) = \mathcal{G}((Y^{-1})^*)$. $\square$

We define X_{even} and X_{odd} by

$$X_{\text{even}} = X^*|_{\text{Image}Y^{\text{even}}}, \quad X_{\text{odd}} = X^*|_{\text{Image}Y^{\text{odd}}}.$$

Proposition 3.11. *If the Jacobi sequence $\{\omega_m\}_{m=1}^\infty$ has property (SC), then*

- X_{even} and X_{odd} are extensions of X ;
- X_{even} and X_{odd} are self-adjoint operators;
- X_{even} and X_{odd} have finite moments of all orders m whose Jacobi sequence is identical to $\{\omega_m\}_{m=1}^\infty$;
- Their Stieltjes transforms $\langle (i - X_{\text{even}})^{-1}\delta_0, \delta_0 \rangle, \langle (i - X_{\text{odd}})^{-1}\delta_0, \delta_0 \rangle$ at i are different.

In the proof, denote the anti-linear isometry $J: \ell_2(\mathbb{N}) \rightarrow \ell_2(\mathbb{N})$ defined by ${}^l(J\eta) = \overline{{}^l\eta}$. Since $JXJ = X$, we have $JX^*J = X^*$. We also have

$$JY^{(m)}J = J(i - X_m)^{-1}J = J(-i - X_m)^{-1}J = ((i - X_m)^*)^{-1} = (Y^{(m)})^*.$$

Taking limits with respect to the strong operator topology, we have

$$JY^{\text{even}}J = (Y^{\text{even}})^*, \quad JY^{\text{odd}}J = (Y^{\text{odd}})^*.$$

Proof. Note that the operator $Y^{\text{even}}: \ell_2(\mathbb{N}) \rightarrow \text{Image}Y^{\text{even}}$ is injective by Lemma 3.8 and that the operator $(Y^{\text{even}})^* = JY^{\text{even}}J$ is also injective. Since $i - X_{\text{even}}$ is equal to $(Y^{\text{even}})^{-1}$, Lemma 3.10 yields that

$$(i - X_{\text{even}})^* = ((Y^{\text{even}})^*)^{-1}.$$

By Lemma 3.8, we have $(-i - X^*)(Y^{\text{even}})^* = J(i - X^*)Y^{\text{even}}J = \text{id}_{\ell_2(\mathbb{N})}$. This means that

$$((Y^{\text{even}})^*)^{-1} = (-i - X^*)|_{\text{Image}(Y^{\text{even}})^*}.$$

By Lemma 3.9, we have

$$(-i - X^*)|_{\text{Image}(Y^{\text{even}})^*} = (-i - X^*)|_{\text{Image}Y^{\text{even}}} = -i - X_{\text{even}}.$$

It follows that $(i - X_{\text{even}})^* = -i - X_{\text{even}}$. We conclude that X_{even} is self-adjoint. The same proof works for X_{odd} .

By taking adjoints of the operators $X_{\text{even}} \subset X^*$, we obtain the inclusion $X \subset \overline{X} \subset X_{\text{even}}$. We also have $X \subset X_{\text{odd}}$. These operators have the same moments with respect to the state $\langle \cdot, \delta_0 \rangle$. The Stieltjes transform of X_{even} at i is equal to

$$\langle Y^{\text{even}} \delta_0, \delta_0 \rangle = \lim_m \langle Y^{(2m)} \delta_0, \delta_0 \rangle = \lim_m iG_{2m}(i) = iG_{\text{even}}.$$

The Stieltjes transform of X_{odd} at i is equal to iG_{odd} . Since property (SC) holds, these values are different (see Proposition 2.6). $\square$

Theorem 3.12. *If the Jacobi sequence $\{\omega_n\}_{n=0}^\infty$ has property (SC), then there exist two probability measures μ_{even} and μ_{odd} such that*

- *These measures μ_{even} and μ_{odd} have finite moments of all orders n whose Jacobi sequence is identical to ω_n ;*
- *These measures are symmetric under the reflexion $\mathbb{R} \ni x \mapsto -x \in \mathbb{R}$;*
- *Their Stieltjes transforms at i are different. That is,*

$$\int_{x \in \mathbb{R}} \frac{1}{i - x} d\mu_{\text{even}} \neq \int_{x \in \mathbb{R}} \frac{1}{i - x} d\mu_{\text{odd}}.$$

Proof. Let $\int_{\mathbb{R}} x dE_{\text{even}}$ be the spectral decomposition of X_{even} and let $\int_{\mathbb{R}} x dE_{\text{odd}}$ be the spectral decomposition of X_{odd} . Define a probability measure μ_{even} by $\langle E_{\text{even}}(\cdot) \delta_0, \delta_0 \rangle$ and define μ_{odd} by $\langle E_{\text{odd}}(\cdot) \delta_0, \delta_0 \rangle$. The measures μ_{even} and μ_{odd} satisfy the first and the third assertions.

Let $P(z, \bar{z}) \in \mathbb{C}[z, \bar{z}]$ be an arbitrary commutative polynomial. Let μ_m denote the probability measure corresponding to X_m and the state $\langle \cdot, \delta_0 \rangle$. Define $p(x)$ by $P((i - x)^{-1}, (-i - x)^{-1})$. Note that $p(x)$ is equal to some polynomial function on the support of μ_m . Since the moments of odd orders $\langle X_m^{2n+1} \delta_0, \delta_0 \rangle$ are zero, we have

$$\int_{\mathbb{R}} p(x) d\mu_m = \int_{\mathbb{R}} p(-x) d\mu_m.$$

The left hand side is equal to

$$\int_{\mathbb{R}} P((i - x)^{-1}, (-i - x)^{-1}) d\mu_m = \langle P(Y^{(m)}, (Y^{(m)})^*) \delta_0, \delta_0 \rangle.$$

The right hand side is equal to

$$\begin{aligned} \int_{\mathbb{R}} P((i + x)^{-1}, (-i + x)^{-1}) d\mu_m &= \int_{\mathbb{R}} P(-(-i - x)^{-1}, -(i - x)^{-1}) d\mu_m \\ &= \langle P(-(Y^{(m)})^*, -Y^{(m)}) \delta_0, \delta_0 \rangle. \end{aligned}$$

Thus we obtain $\langle P(Y^{(m)}, (Y^{(m)})^*) \delta_0, \delta_0 \rangle = \langle P(-(Y^{(m)})^*, -Y^{(m)}) \delta_0, \delta_0 \rangle$. Taking a limit with respect to the strong operator topology, we obtain

$$\langle P(Y^{\text{even}}, (Y^{\text{even}})^*) \delta_0, \delta_0 \rangle = \langle P(-(Y^{\text{even}})^*, -Y^{\text{even}}) \delta_0, \delta_0 \rangle,$$

and hence

$$\begin{aligned} \int_{\mathbb{R}} P((i-x)^{-1}, (-i-x)^{-1}) d\mu_{\text{even}} &= \int_{\mathbb{R}} P((i+x)^{-1}, (-i+x)^{-1}) d\mu_{\text{even}}, \\ \int_{\mathbb{R}} p(x) d\mu_{\text{even}} &= \int_{\mathbb{R}} p(-x) d\mu_{\text{even}}. \end{aligned}$$

Since the polynomials $\{P((i-x)^{-1}, (-i-x)^{-1})\}$ form a dense subspace of the C^* -algebra $C_0(\mathbb{R})$, we have $d\mu_{\text{even}}(x) = d\mu_{\text{even}}(-x)$. The measure μ_{odd} is also symmetric. $\square$

4. EXAMPLES OF INFINITE JACOBI SEQUENCES WITH PROPERTY (SC)

When we seek examples of property (SC), the following are useful.

Lemma 4.1. $C_n < \frac{1}{\omega_n} + \frac{\omega_{n-1}}{\omega_n} C_{n-2} \quad n \geq 2.$

$$\begin{aligned} C_{2n} &< \frac{1}{\omega_{2n}} + \frac{\omega_{2n-1}}{\omega_{2n}} \frac{1}{\omega_{2n-2}} + \cdots + \left(\frac{\omega_{2n-1}}{\omega_{2n}} \cdots \frac{\omega_3}{\omega_4} \frac{1}{\omega_2} \right) + \left(\frac{\omega_{2n-1}}{\omega_{2n}} \cdots \frac{\omega_3}{\omega_4} \frac{\omega_1}{\omega_2} \right), \\ C_{2n+1} &< \frac{1}{\omega_{2n+1}} + \frac{\omega_{2n}}{\omega_{2n+1}} \frac{1}{\omega_{2n-1}} + \cdots + \left(\frac{\omega_{2n}}{\omega_{2n+1}} \cdots \frac{\omega_2}{\omega_3} \frac{1}{\omega_1} \right), \quad n \geq 1. \end{aligned}$$

Proof. By the definition of C_n , we have

$$C_n = \frac{1}{1 + \frac{\omega_n}{1 + \omega_{n-1} C_{n-2}}} < \frac{1 + \omega_{n-1} C_{n-2}}{\omega_n} = \frac{1}{\omega_n} + \frac{\omega_{n-1}}{\omega_n} C_{n-2}.$$

Applying this inequality to C_{n-2} , we have

$$\begin{aligned} C_n &< \frac{1}{\omega_n} + \frac{\omega_{n-1}}{\omega_n} C_{n-2} \\ &< \frac{1}{\omega_n} + \frac{\omega_{n-1}}{\omega_n} \left(\frac{1}{\omega_{n-2}} + \frac{\omega_{n-3}}{\omega_{n-2}} C_{n-4} \right) \\ &= \frac{1}{\omega_n} + \frac{\omega_{n-1}}{\omega_n} \frac{1}{\omega_{n-2}} + \frac{\omega_{n-1}}{\omega_n} \frac{\omega_{n-3}}{\omega_{n-2}} C_{n-4}. \end{aligned}$$

Repeating this procedure, we obtain the second and the third inequalities. $\square$

4.1. Jacobi sequences with high growth rates. We study the case that the infinite Jacobi sequence $\{\omega_n\}$ satisfies the following condition:

(*) There exist $0 < \alpha < 1$ and $n_0 \in \mathbb{N}$ such that $\omega_n < \alpha \omega_{n+1}$ for $n \geq n_0$.

This condition implies that there exists a constant $K > 0$ such that $1/\omega_n < K\alpha^n$.

Theorem 4.2. *The sequence $\{\omega_n\}$ with condition (*) has property (SC). As a consequence, the corresponding probability measure is not unique.*

Proof. Define an even number $2m_0$ by n_0 or $n_0 - 1$. By the first inequality in Lemma 4.1, for $n \geq n_0 + 1$, we have $C_n < K\alpha^n + \alpha C_{n-2}$. Applying this inequality many times, we obtain for $2n > n_0 + 1$,

$$\begin{aligned} & C_{2n} \\ & < K\alpha^{2n} + \alpha K\alpha^{2n-2} + \dots + \alpha^{(2n-2m_0)/2-1} K\alpha^{2m_0+2} + \alpha^{(2n-2m_0)/2} C_{2m_0} \\ & < \max\{K, C_{2m_0}\} (\alpha^{2n} + \alpha^{2n-1} + \dots + \alpha^{n+m_0+1} + \alpha^{n+m_0}) \\ & < \max\{K, C_{2m_0}\} \cdot \alpha^n / (1 - \alpha). \end{aligned}$$

We obtain for $2n + 1 > n_0 + 1$,

$$\begin{aligned} & C_{2n+1} \\ & < K\alpha^{2n+1} + \dots + \alpha^{(2n-2m_0)/2-1} K\alpha^{2m_0+3} + \alpha^{(2n-2m_0)/2} C_{2m_0+1} \\ & < \max\{K, C_{2m_0+1}\} (\alpha^{2n+1} + \alpha^{2n} + \dots + \alpha^{n+m_0+1} + \alpha^{n+m_0+1}) \\ & < \max\{K, C_{2m_0+1}\} \cdot \alpha^n / (1 - \alpha). \end{aligned}$$

It follows that $\sum_{n=1}^{\infty} C_n = \sum_{n=1}^{n_0+1} C_n + \sum_{n=n_0+2}^{\infty} C_n < \infty$. $\square$

4.2. Jacobi sequences for q -Gaussians. The Jacobi sequence of the Gaussian measure $e^{-x^2/2} dx / \sqrt{2\pi}$ is the sequence $\{n = [n]_1\}$ of integers. In this subsection, we study the following deformations of the sequence:

- $\omega_n = [n]_q = 1 + q + q^2 + \dots + q^{n-1} = \frac{q^n - 1}{q - 1}, \quad q > -1,$
- $\omega_n = (-1)^{n-1} \frac{q^n - 1}{q - 1}, \quad q < -1.$

The number $[n]_q$ is called a q -deformed integer. Their corresponding probability measures are often called “ q -Gaussians.” The study of q -Gaussians are motivated by the study of q -canonical commutation relations and q -canonical anti-commutation relations. See [Bož07] and references therein.

Theorem 4.3. *The moment problem for q -Gaussians ($q \in \mathbb{R}, q \neq -1$) are indeterminate if and only if $|q| > 1$.*

Proof. If $|q| > 1$, the Jacobi sequence satisfies condition (*), since

$$\lim_n \frac{\omega_{n+1}}{\omega_n} = \lim_n \left| \frac{q^{n+1} - 1}{q^n - 1} \right| = |q| > 1.$$

By Theorem 4.2, the Jacobi sequence has property (SC). By Theorem 3.12, there are more than two symmetric probability measures corresponding to ω_n . In the case that $-1 < q \leq 1$, we have $[q]_n \leq n$. Carleman’s condition states that if $\{\omega_n\}$ satisfies the condition $\sum_n 1/\sqrt{\omega_n} = +\infty$, then the corresponding probability measure is unique. Theorem 5.1 proved later also yields the results. $\square$

Remark 4.4. For $-1 < q \leq 1$, the theorem is well-known. For $q > 1$, Ismail–Masson [IM94] has already shown the indeterminacy.

4.3. Power of integers. We are going to consider the case of $\{\omega_n = n^p\}_{n=1}^\infty$.

Lemma 4.5. *If $\omega_n = n^p$ and $p \geq 0$, then*

$$\frac{\omega_n}{\omega_{n+1}} < \sqrt{\frac{\omega_n}{\omega_{n+2}}}, \quad \frac{\sqrt{\omega_n}}{\omega_{n-1}} < 2^p \frac{1}{\sqrt{\omega_n}} \quad (n \geq 2).$$

Proof. By the inequality $n/(n+1) < (n+1)/(n+2)$, we have

$$\sqrt{\omega_n/\omega_{n+1}} < \sqrt{\omega_{n+1}/\omega_{n+2}}.$$

Thus we obtain the following inequality:

$$\frac{\omega_n}{\omega_{n+1}} = \sqrt{\frac{\omega_n}{\omega_{n+1}}} \sqrt{\frac{\omega_n}{\omega_{n+1}}} < \sqrt{\frac{\omega_n}{\omega_{n+1}}} \sqrt{\frac{\omega_{n+1}}{\omega_{n+2}}} = \sqrt{\frac{\omega_n}{\omega_{n+2}}}.$$

The second inequality is due to the following computation:

$$\frac{\sqrt{\omega_n}}{\omega_{n-1}} = \frac{\omega_n}{\omega_{n-1}} \frac{1}{\sqrt{\omega_n}} < \left(\frac{n}{n-1}\right)^p \frac{1}{\sqrt{\omega_n}} < 2^p \frac{1}{\sqrt{\omega_n}}.$$

□

Proposition 4.6. *For every $p > 2$, the Jacobi sequence $\{n^p\}_{n=1}^\infty$ has property (SC). As a consequence, there exist more than two symmetric probability measures whose Jacobi sequence is $\{n^p\}_{n=1}^\infty$.*

Proof. By Lemma 4.5, we have

$$\begin{aligned} \frac{\omega_{n-1}}{\omega_n} \frac{\omega_{n-3}}{\omega_{n-2}} \dots \frac{\omega_l}{\omega_{l+1}} \frac{1}{\omega_{l-1}} &< \sqrt{\frac{\omega_{n-1}}{\omega_{n+1}}} \sqrt{\frac{\omega_{n-3}}{\omega_{n-1}}} \dots \sqrt{\frac{\omega_l}{\omega_{l+2}}} \frac{1}{\omega_{l-1}} \\ &= \frac{1}{\sqrt{\omega_{n+1}}} \frac{\sqrt{\omega_l}}{\omega_{l-1}} \\ &< 2^p \frac{1}{\sqrt{\omega_{n+1}}} \frac{1}{\sqrt{\omega_l}}. \end{aligned}$$

According to Lemma 4.1, the following inequalities hold:

$$\begin{aligned} C_{2n} &< 2^p \frac{1}{\sqrt{\omega_{2n+1}}} \left(\frac{1}{\sqrt{\omega_{2n+1}}} + \frac{1}{\sqrt{\omega_{2n-1}}} + \dots + \frac{1}{\sqrt{\omega_3}} + \frac{1}{\sqrt{\omega_1}} \right), \\ C_{2n+1} &< 2^p \frac{1}{\sqrt{\omega_{2n+2}}} \left(\frac{1}{\sqrt{\omega_{2n+2}}} + \frac{1}{\sqrt{\omega_{2n}}} + \dots + \frac{1}{\sqrt{\omega_4}} + \frac{1}{\sqrt{\omega_2}} \right). \end{aligned}$$

By the inequality

$$\sum_{n=1}^{\infty} C_n < \sum_{n=1}^{\infty} 2^p \frac{1}{\sqrt{\omega_{n+1}}} \sum_{l=1}^{\infty} \frac{1}{\sqrt{\omega_l}} < 2^p \left(\sum_{n=1}^{\infty} n^{-p/2} \right)^2,$$

we conclude that $\{\omega_n = n^p\}_{n=1}^\infty$ have property (SC). □

By the same discussion in the proof of Theorem 4.3, we obtain the following:

Theorem 4.7. *Let p be a positive number. The probability measure corresponding to the Jacobi sequence $\{n^p\}$ is unique if and only if $p \leq 2$.*

Applying the classical result on orthogonal polynomials (due to Meixner [Mei]), it is easy to show that the sequence is corresponding to the probability measure $\frac{1}{e^{\pi x/2} + e^{-\pi x/2}} dx$, which is now called “hyperbolic secant distribution.” The equation (5, 4) of [Mei] gives a sequence of polynomials $\{P_n(x)\}_{n=-1,0,\dots}$ satisfying $xP_n = P_{n+1} + n^2 P_{n-1}$ in the case that $l_1 = 0$, $\lambda = 0$, $k_2 = -1$ and $\kappa = -1$. The last paragraph of [Mei] proves that the three-term recurrence relation is realized by the probability measure $d\psi$. The measure is a scalar multiple of

$$\Gamma\left(\frac{ix+1}{2}\right) \Gamma\left(\frac{-ix+1}{2}\right) dx = \frac{\pi}{\sin((ix+1)\pi/2)} dx = \frac{\pi}{\cos(ix\pi/2)} dx.$$

Corollary 4.7 means that for the case of $\omega_n = n^p$ the hyperbolic secant distribution is “the last probability measure” which is uniquely determined by the moment sequence.

5. DETERMINATE MOMENT PROBLEM AND PROPERTY (SC)

Theorem 5.1. *Let $\{M_n\}_{n=0}^\infty$ be a sequence of moments. Suppose that $M_n = 0$ for every odd number n . Let $\{\omega_n\}_{n=1}^\infty$ denote the corresponding Jacobi sequence. Suppose that the Jacobi sequence $\{\omega_n\}_{n=1}^\infty$ is of infinite type. Then the following conditions are equivalent:*

- (1) *The sequence $\{\omega_n\}_{n=1}^\infty$ has property (SC);*
- (2) *A probability measure μ satisfying $M_n = \int_{x \in \mathbb{R}} x^n d\mu$ is not unique.*

We have already shown that if condition (1) holds, then there exist two probability measures μ_{even} and μ_{odd} which are distinguished by the Stieltjes transforms at i . We prove that condition (2) implies (1).

Proof. Suppose that a probability measure μ satisfying $M_n = \int_{x \in \mathbb{R}} x^n d\mu$ is not unique. By Theorem 2 of Simon [Sim98], the symmetric operator X is not essentially self-adjoint. Then the kernel $\ker(i - X^*)$ or $\ker(-i - X^*)$ contains a non-zero vector ξ . Replacing i with $-i$, if necessary, we may assume that $\ker(i - X^*)$ contains a non-zero vector $Z = ({}^n Z)_{n=0}^\infty$. Then for every k , we have $\langle Z, (-i - X)\delta_k \rangle = 0$ and hence

$$i({}^0 Z) - \sqrt{\omega_1}({}^1 Z) = 0, \quad -\sqrt{\omega_k}({}^{k-1} Z) + i({}^k Z) - \sqrt{\omega_{k+1}}({}^{k+1} Z) = 0$$

By induction, the n -th entry of the vector can be expressed by the sequence B_n as

$${}^n Z = {}^0 Z \frac{B_{n-1} i^n}{\sqrt{\omega_1 \cdots \omega_n}}, \quad n \geq 1.$$

For $n \geq 2$, by the equality $C_n = B_{n-1}/B_n$, we have

$$\frac{({}^n Z)^2}{({}^{n-1} Z)^2} = \frac{B_{n-1}^2}{\omega_n B_{n-2}^2} = \frac{B_{n-1}^2}{(B_n - B_{n-1})B_{n-2}} = \frac{C_n}{(1 - C_n)C_{n-1}}.$$

Thus we have

$$({}^n Z)^2 = ({}^0 Z)^2 \prod_{l=1}^n \frac{C_l}{(1 - C_l)C_{l-1}} = ({}^0 Z)^2 \frac{C_n}{\prod_{l=1}^n (1 - C_l)} \geq ({}^0 Z)^2 C_n.$$

By the inequality $\sum_{n=2}^{\infty} C_n \leq \|Z\|_2^2 / ({}^0Z)^2$, we conclude that property (SC) holds. $\square$

Acknowledgment. The authors are grateful to Prof. M. Bożejko for many suggestions and comments to refine the drafts.

REFERENCES

- [Akh65] N. I. Akhiezer, *The classical moment problem and some related questions in analysis*, Translated by N. Kemmer, Hafner Publishing Co., New York, 1965.
- [Boż07] M. Bożejko, *Remarks on q -CCR relations for $|q| > 1$* , Banach Center Publ. **78** (2007), 59–76.
- [Ham20a] H. Hamburger, *Über eine Erweiterung des Stieltjesschen Momentenproblems*, Math. Ann. **81** (1920), no. 2-4, 235–319.
- [Ham20b] ———, *Über eine Erweiterung des Stieltjesschen Momentenproblems*, Math. Ann. **82** (1920), no. 1-2, 120–164.
- [Ham21] ———, *Über eine Erweiterung des Stieltjesschen Momentenproblems*, Math. Ann. **82** (1921), no. 3-4, 168–187.
- [HO07] A. Hora and N. Obata, *Quantum probability and spectral analysis of graphs*, Theoretical and Mathematical Physics, Springer, Berlin, 2007.
- [IM94] M. E. H. Ismail and D. R. Masson, *q -Hermite polynomials, biorthogonal rational functions, and q -beta integrals*, Trans. Amer. Math. Soc. **346** (1994), no. 1, 63–116.
- [Mei] J. Meixner, *Orthogonale Polynomsysteme Mit Einer Besonderen Gestalt Der Erzeugenden Funktion*, J. London Math. Soc. **S1-9**, no. 1, 6–13.
- [Sim98] B. Simon, *The classical moment problem as a self-adjoint finite difference operator*, Adv. Math. **137** (1998), no. 1, 82–203.
- [ST43] J. A. Shohat and J. D. Tamarkin, *The Problem of Moments*, American Mathematical Society Mathematical surveys, vol. I, American Mathematical Society, New York, 1943.
- [Sti94] T.-J. Stieltjes, *Recherches sur les fractions continues*, Ann. Fac. Sci. Toulouse Sci. Math. Sci. Phys. **8** (1894), no. 4, J1–J122.

HAYATO SAIGO, NAGAHAMA INSTITUTE OF BIO-SCIENCE AND TECHNOLOGY, NAGAHAMA 526-0829, JAPAN

E-mail address: h_saigoh@nagahama-i-bio.ac.jp

HIROKI SAKO, SCHOOL OF SCIENCE, TOKAI UNIVERSITY, KANAGAWA, 259-1292 JAPAN

E-mail address: hiroki@tokai-u.jp